



King's Research Portal

DOI:

[10.1136/thoraxjnl-2017-210422](https://doi.org/10.1136/thoraxjnl-2017-210422)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Shankar-Hari, M., & McAuley, D. F. (2017). Divide and conquer: identifying acute respiratory distress syndrome subphenotypes. *Thorax*, 72(10), 867-869. <https://doi.org/10.1136/thoraxjnl-2017-210422>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Divide and conquer: Identifying Acute Respiratory Distress Syndrome sub-phenotypes

Manu Shankar-Hari^{1,2} and Daniel F. McAuley^{3,4#}

¹Department of Critical Care Medicine, Guy's and St Thomas' NHS Foundation Trust, Westminster Bridge Rd, London SE17EH, United Kingdom; ²Division of Infection and Immunity, King's College London, United Kingdom; ³Centre for Experimental Medicine, Wellcome-Wolfson Institute for Experimental Medicine, Queen's University of Belfast, Belfast, Northern Ireland; ⁴Regional Intensive Care Unit, Royal Victoria Hospital, Grosvenor Road, Belfast, Northern Ireland

Manu Shankar-Hari^{1,2} M.Sc., Ph.D.

email: manu.shankar-hari@kcl.ac.uk

Twitter: @msh_manu

#Corresponding author

Professor Daniel F McAuley

³Centre for Experimental Medicine, Wellcome-Wolfson Institute for Experimental Medicine, Queen's University of Belfast, Belfast, Northern Ireland; ⁴Regional Intensive Care Unit, Royal Victoria Hospital, Grosvenor Road, Belfast, Northern Ireland

email: d.f.mcauley@qub.ac.uk

Tel: 0044 28 90635794

Word count = 1025 (excluding Table)

References = 13

Figures and tables = 1 table

Key words: ARDS, Humans, Phenotypes, Enrichment; Stratified Medicine

The acute respiratory distress syndrome (ARDS) definition identifies patients with acute onset hypoxaemia and respiratory failure, who have bilateral opacities on chest radiograph that are not fully explained by cardiac failure or fluid overload¹. ARDS is a common illness that accounts for approximately 10% of critical care admissions and 20% of patients requiring mechanical ventilation². The hospital mortality in patients with ARDS remains high, increasing from approximately 35% for those with mild disease to 46% for those with severe ARDS². This high mortality has remained relatively unchanged in the last 20 years³. To date, despite decades of research, there is no pharmacological treatment that can modify the underlying biological mechanisms implicated in ARDS and improve patient outcomes⁴. Within ARDS populations there is substantial biological and outcome heterogeneity, with observed differences in dominant pathogenic mechanisms, treatment responses and outcomes⁵⁻⁷. Identifying ARDS sub-phenotypes based on biological characteristics mechanistically linked to specific therapies irrespective of the baseline risk of outcome, is the conceptual definition of predictive enrichment^{7,8}. The identification of such ARDS sub-phenotypes will enable improved trial design in ARDS by selecting patients based on responder characteristics to therapeutic interventions, hopefully resulting in improved outcomes⁶.

In this issue of Thorax, Bos et al report an excellent cohort study in 700 ARDS patients, testing the hypothesis that ARDS sub-groups exist due to difference in biological characteristics⁹. In this retrospective analysis of a prospectively collected cohort, 20 biomarkers were selected to represent inflammation, coagulation and endothelial activation, as hallmarks of ARDS biology⁶. The dataset was divided into a training cohort (n=454 patients) and validation cohort (n=246 patients), based on the study recruitment period. Cluster analysis was used to identify homogenous ARDS sub-phenotypes in the training cohort¹⁰. The most predictive biomarkers were then confirmed in the validation cohort. These biological clusters were then linked to clinical and outcome characteristics of ARDS patients to derive clinical sub-phenotypes, namely *reactive* and *uninflamed*. These two clinical ARDS sub-phenotypes differed in terms of illness severity and critical care mortality, with the reactive group having a greater risk of death.

A key question for the reader is whether these associations are spurious or indirect or causal?¹¹ Cluster analysis methods generate different results dependent on the variables chosen for identifying similarities between patients and the method of clustering¹⁰. Bos et al chose biomarker characteristics as the variables on which the groups should be similar, and used Ward's method of

agglomerative hierarchical clustering to identify two potentially generalizable ARDS clusters. Hierarchical clustering is a commonly used iterative method to identify homogenous groups or clusters based on specific characteristics. In the paper by Bos et al, the goal was to identify ARDS patients with similar biomarker profiles, from a heterogeneous ARDS cohort. The basic algorithm starts with assigning each ARDS patient a 'value' based on their individual biomarker profile. Then patients with similar 'values' are grouped together to form clusters. The underlying principle is that ARDS patients *within* each cluster will have similar biomarker profiles and that *between* clusters biomarker profiles will be different. Depending on the parameters specified, the same dataset can result in potentially different results with different clustering algorithms and there are no universally agreed optimal rule(s) for clustering¹⁰. Another potential limitation is that only patients with data on all chosen biomarkers were included and missing data in clinical variables were imputed, which has the potential for selection and information bias. The blood sampling window for biomarker measurement in this cohort was wide and drawn either on the day of ARDS diagnosis or the day before or the day after, challenging the time-based arguments for a causal relationship. Despite these challenges, Bos et al provide important data with strong associations that are consistent with our current knowledge, have biological plausibility and external validity.

Calfee and colleagues have led the field in defining ARDS sub-phenotypes. Using latent class analysis (LCA) of clinical and biomarker data from patients enrolled in ARDS randomised controlled trials, Calfee *et al* have originally identified two ARDS sub-phenotypes^{12 13}. The *reactive* sub-phenotype identified in this study shares many of the features of the *hyperinflammatory* ARDS sub-phenotype reported previously^{12 13}, although the proportion of patients in the *reactive* group is much higher than the *hyperinflammatory* sub-phenotype. This suggests that the *hyperinflammatory* and *reactive* groups may represent a similar sub-phenotype, although this is unproven. The findings from this study are significant in that they have identified comparable sub-phenotypes in an observational cohort of patients with ARDS using a different analytic approach. Whilst Calfee et al identified these ARDS sub phenotypes using clinical and biomarker data, Bos et al identified them purely on biomarker data. It would be important to test whether similar sub-phenotypes emerge after harmonising these different study datasets and performing both cluster and latent class analyses. Table 1 provides a comparative summary of these three studies.

Table 1: Summary of studies that report ARDS sub-phenotypes

Parameter	Bos et al ⁹	Calfee C et al ¹³	Famous et al ¹²
Sample size	700	1022	1000
Recruitment period	2011 – 2013	1996 – 2002	2000 - 2005
Study design	Observational cohort	RCT analysed as cohort	RCT analysed as cohort
ARDS P/F criteria	<=300	<300	<300
Blood sampling	Around ARDS diagnosis	At baseline	At baseline
Biomarkers used for deriving sub-phenotypes	Lung epithelial: <i>none</i> Endothelial: E-selectin; P-selectin; ANG1/2 Coagulation: Antithrombin; D-Dimer; tPA; PAI-1; Inflammation: Fractalkine; GM-CSF; ICAM-1; IFN- γ ; IL-1 β ; IL-6; IL-8; IL-10; IL-13; TNF- α ; MMP-8; TIMP-1;	Lung epithelial: SP-D Endothelial: ICAM-1; vWF Coagulation: Protein C; PAI-1 Inflammation: sTNFR-1; IL-6; IL-8	Lung epithelial: SP-D Endothelial: ICAM-1; vWF Coagulation: Protein C; PAI-1 Inflammation: sTNFR-1; IL-6; IL-8
Clinical variables used for deriving sub-phenotypes	None	Age, gender, ethnicity, BMI, Respiratory [#] ; Cardiovascular*; Creatinine; Urine output; Bilirubin; Temperature; Haematocrit; WBC count; Sodium; glucose; Albumin; Platelets; bicarbonate; Aetiology of ARDS [^]	Age, gender, ethnicity, BMI, Respiratory [#] ; Cardiovascular*; Creatinine; Urine output; Bilirubin; Temperature; Haematocrit; WBC count; Sodium; glucose; Albumin; Platelets; bicarbonate; Aetiology of ARDS [^]
Analytical approach to derive ARDS subsets	Cluster analyses based only on biomarker data	Latent class analyses based grouping based on clinical and biomarker data	Latent class analyses based grouping based on clinical and biomarker data
ARDS subset (prevalence %)	Reactive phenotype (58.0%) Vs Uninflamed (42.0%)	Hyper-Inflammatory (29.4%) Vs Phenotype 1 (70.6%)	Hyper-Inflammatory (27.3%) Vs Phenotype 1 (72.7%)
Mortality (%) by ARDS subset	Reactive phenotype = 36.8% Vs Uninflamed = 14.9%	Hyper-Inflammatory = 47.3% Vs Phenotype 1 = 19.4%	Hyper-Inflammatory = 45.0% Vs Phenotype 1 = 22.0%
Discriminant markers between phenotypes	IL-6; IFN-gamma; ANG1/2; PAI-1	IL-6; sTNFR1; Vasopressor use; IL-8; HCO3	IL-8; sTNFR1; Vasopressor use; HCO3; minute ventilation

Legend to Table-1:

Table-1 shows the summary of three recent studies that report ARDS sub-phenotypes. The **Respiratory system variables**[#] included minute ventilation, mean airway pressure, plateau pressure, respiratory rate, tidal volume, positive end-expiratory pressure; partial pressure of carbondioxide (PaCO₂) and PaO₂/FiO₂ ratio. The **Cardiovascular* system** variables include highest heart rate, lowest systolic blood pressure and vasopressor use. The **aetiology of ARDS**[^] was coded as Trauma, Sepsis, Aspiration, Pneumonia or Other. **Abbreviations:** P/F = PaO₂/FiO₂ ratio; ANG1/2 = Angiopoietin 1 and 2; tPA = Tissue plasminogen activator; PAI-1 = plasminogen activator inhibitor-1; GM-CSF = granulocyte-monocyte colony stimulating factor; ICAM-1 = intracellular adhesion molecule-1; IFN- γ = Interferon-gamma; IL-1 β = interleukin-1 beta; Interleukins = IL-6; IL-8; IL-10; IL-13; TNF- α = Tumor necrosis factor-alpha; MMP-8 = matrix metalloproteinase-8; TIMP-1= tissue inhibitor of metalloproteinase-1; SP-D = Surfactant protein-D; vWF = von-Willebrand's Factor; sTNFR-1 = soluble Tumor necrosis factor receptor-1; BMI = body mass index; WBC = white blood cell count;

Several important questions remain unanswered. First, assuming the *hyperinflammatory/reactive* sub-phenotype represents a common sub-phenotype, further work is needed to identify the key discriminant makers to reliably define this ARDS subset. Ideally a minimal data-set of variables could be identified to efficiently achieve this. Second, although it remains unknown if ARDS sub-phenotypes respond differently to pharmacotherapies, an important aspect of in developing pharmacotherapies targeted at the *hyperinflammatory/reactive* sub-phenotype will be to determine the stability of the ARDS subgroup over time. This is important to determine the therapeutic window to intervene with a therapy targeted at this sub-phenotype. In addition, it would be important to define if and how moving from this sub-phenotype to an *uninflamed* phenotype represents therapeutic success or failure to guide ongoing treatment. Third, development of point-of-care assays along with algorithms to define these ARDS sub-phenotypes at the bedside in real-time is essential to enable this information to inform clinical trials targeting these sub-phenotypes.

In summary, ARDS continues to be a clinical and research challenge in terms of developing pharmacological therapies. Bos et al provide intriguing data that highlights the need for further work to identify ARDS subsets with defined treatable traits. These sub-phenotypes should be based on modifiable biological characteristics linked to both the risk of poor outcomes and response to the tested treatment. This will enable personalised care of patients with ARDS.

Acknowledgments: MS-H is supported by the National Institute for Health Research Clinician Scientist Award (NIHR-CS-2016-16-011). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the UK National Institute for Health Research or the Department of Health.

Ethics approval and consent to participate: Not applicable

Availability of data and material: Not applicable

Competing interests: The author declares that they have no competing interests

Funding: Not applicable

Authors' contributions: MSH wrote the first draft. MSH and DM critically revised the manuscript for important intellectual content and agreed the final submitted version of the manuscript.

References

1. Force ADT, Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307(23):2526-33. doi: 10.1001/jama.2012.5669 [published Online First: 2012/07/17]
2. Bellani G, Laffey JG, Pham T, et al. Epidemiology, Patterns of Care, and Mortality for Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries. *JAMA* 2016;315(8):788-800. doi: 10.1001/jama.2016.0291
3. Phua J, Badia JR, Adhikari NK, et al. Has mortality from acute respiratory distress syndrome decreased over time?: A systematic review. *Am J Respir Crit Care Med* 2009;179(3):220-7. doi: 10.1164/rccm.200805-722OC
4. Boyle AJ, Sweeney RM, McAuley DF. Pharmacological treatments in ARDS; a state-of-the-art update. *BMC Med* 2013;11(1):166. doi: 10.1186/1741-7015-11-166
5. Pham T, Rubenfeld GD. Fifty Years of Research in ARDS. The Epidemiology of ARDS: A Fiftieth Birthday Review. *Am J Respir Crit Care Med* 2017 doi: 10.1164/rccm.201609-1773CP [published Online First: 2017/02/06]
6. Sweeney RM, McAuley DF. Acute respiratory distress syndrome. *Lancet* 2016;388(10058):2416-30. doi: 10.1016/S0140-6736(16)00578-X
7. Shankar-Hari M, Rubenfeld GD. The use of enrichment to reduce statistically indeterminate or negative trials in critical care. *Anaesthesia* 2017:n/a-n/a. doi: 10.1111/anae.13870
8. Prescott HC, Calfee CS, Thompson BT, et al. Toward Smarter Lumping and Smarter Splitting: Rethinking Strategies for Sepsis and Acute Respiratory Distress Syndrome Clinical Trial Design. *Am J Respir Crit Care Med* 2016;194(2):147-55. doi: 10.1164/rccm.201512-2544CP [published Online First: 2016/06/01]
9. Bos L et al. Identification and validation of distinct biological phenotypes in patients with acute respiratory distress syndrome by cluster analysis. *Thorax* (2017) In press.
10. Frades I, Matthiesen R. Overview on techniques in cluster analysis. *Methods Mol Biol* 2010;593:81-107. doi: 10.1007/978-1-60327-194-3_5 [published Online First: 2009/12/04]
11. Grimes DA, Schulz KF. Bias and causal associations in observational research. *Lancet* 2002;359(9302):248-52. doi: 10.1016/S0140-6736(02)07451-2
12. Famous KR, Delucchi K, Ware LB, et al. Acute Respiratory Distress Syndrome Subphenotypes Respond Differently to Randomized Fluid Management Strategy. *Am J Respir Crit Care Med* 2017;195(3):331-38. doi: 10.1164/rccm.201603-0645OC
13. Calfee CS, Delucchi K, Parsons PE, et al. Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. *The Lancet Respiratory medicine* 2014;2(8):611-20. doi: 10.1016/S2213-2600(14)70097-9